

# JCU ePrints

This file is part of the following reference:

**Moore, Leslie Allan (2010) *Niche differentiation, rarity, and commonness in the sympatric Australian white-tailed rats: *Uromys caudimaculatus* and *Uromys hadrourus**. PhD thesis, James Cook University.**

Access to this file is available from:

<http://eprints.jcu.edu.au/17434>



## REFERENCES

- Akcakaya, H., Burgman, M., and L. Ginzburg (1999). Applied population ecology. 2nd edition. Sinauer, Sunderland, Massachusetts.
- Alatalo, R. (1981). Problems in the measurement of evenness in ecology. *Oikos* 37:199–204.
- Albrecht, M., and N. Gotelli (2001). Spatial and temporal niche partitioning in grassland ants. *Oecologia* (2001) 126:134–141.
- Amstrup, S., McDonald, T., and B. Manly (eds.). 2005. Handbook of Capture-Recapture Analysis. Princeton University Press: Princeton, New Jersey.
- Aplin, K. (2006). Ten million years of rodent evolution in Australasia: Phylogenetic evidence and a speculative historical biogeography. In: Merrick, J., Archer, M., Hickey, G., and Lee, M. (eds). Evolution and biogeography of Australasian vertebrates. Auscipub Pty Ltd. NSW.
- Aplin, K., Baverstock, P., and S. Donnellan (1993). Albumin immunological evidence for the time and mode of origin of the New Guinea terrestrial mammal fauna. *Science in New Guinea*. Vol. 19, 131-144.
- Aplin, K., Brown, P., Jacob, J., Krebs, C., and G. Singleton (2003). Field methods for rodent studies in Asia and the Indo-Pacific. (ACIAR Monograph No. 100).
- Arita, H., Robinson, J., and K. Redford (1990). Rarity in Neotropical forest mammals and its ecological correlates. *Conservation Biology*, 4, 181-192.
- Arita (1993). Rarity in Neotropical Bats: Correlations with Phylogeny, Diet, and Body Mass. *Ecological Applications* (3), 506-517.
- Armstrong, R., and R. McGehee (1980). Competitive exclusion. *American Naturalist* Vol. 115, 151.
- Balston, J. (2009). Impacts of climate variability and climate change on the Wet Tropics of north-eastern Australia. In: Living in a dynamic tropical forest landscape. Eds Nigel Stork and Stephen Turton: Blackwell Publishing.

- Barber, S., and L. Mech. (2002). A Critique of Wildlife Radio-tracking and its Use in National Parks. *A Report to the U.S. National Park Service*.
- Baverstock, P., Watts, C., and J. Hogarth (1977a). Chromosome evolution in Australian rodents. I. The Pseudomyinae, the Hydrominae, and the *Uromys/Melomys* group. *Chromosoma*, Vol. 61, 95-125.
- Baverstock, P., Watts, C., Hogarth, J., Robinson, J. T., and J. Robinson (1977b). Chromosome evolution in Australian rodents. II. *The Rattus group*. *Chromosoma*, Vol. 61, 227-241.
- Baverstock, P., Watts, C., Adams, M., and S. Cole (1981). Genetical relationships among Australian rodents (Muridea). *Australian Journal of Zoology*. Vol. 29, 289-303.
- Begon, M. (1983). Abuses of mathematical techniques in ecology: applications of Jolly's capture-recapture method. *Oikos* 40, 155-158.
- Begon, M., Harper, L., and C. Townsend (1996). *Ecology: Individuals, Populations, and Communities*, 3rd edition. Blackwell Science Ltd. Cambridge, MA.
- Bell, G. (2001). Neutral Macroecology. *Science* (293) 2413 – 2418.
- Benayas, R., Scheiner, Sánchez-Colomer, M., and C. Levassor. (1999). Commonness and rarity: theory and application of a new model to Mediterranean montane grasslands. *Conservation Ecology* [online] 3(1).
- Betancourt, J., Van Devender, T., and M. Rose (1986). Comparison of plant microfossils in woodrat (*Neotoma* sp.) and porcupine (*Erethizon dorsatum*) middens from the western United States. *J.Mamm.* 67(2): 266-273.
- Blackburn, T., and K. Gaston. (1996) A sideways look at patterns in species richness, or why there are so few species outside the tropics. *Biodiversity Letters* 3: 44-53.

- Blackburn, T., and K. Gaston (1997). Who is rare? Artefacts and complexities of rarity determination. In: W. Kunin and K. Gaston (eds), *The Biology of rarity, Population and Community Biology Series*, Chapman and Hall, London, pp. 48-58.
- Blackburn, T., Brown, V., and Doube (1993). The relationship between body size and abundance in natural animal assemblages. *Journal of Animal Ecology*, 62, 519-528.
- Blackburn, T., Gates, S., Lawton, J. and J. Greenwood (1994). Relations between body size, abundance, and taxonomy of birds wintering in Britain and Ireland. *Philosophical Transactions of the Royal Society, London, B*, 343, 135-144.
- Bohlin, T., Dellefors, C., Faremo, U., and A. Johlander (1994). The energetic equivalence hypothesis and the relation between population density and body size in stream-living salmonids. *Am. Nat.* 143: 478-493.
- Boutin, S. (1990). Food supplementation experiments with terrestrial vertebrates: patterns, problems, and the future. *Canadian Journal of Zoology*, 68, 203-220.
- Bowman, J., Jaeger, J., and L. Fahrig (2002). Dispersal distance of mammals is proportional to home range size. *Ecology*, 83, 2049–2055.
- Brandle, M., Ohlschlager, S. and R. Brandl (2002). Range sizes in butterflies: correlation across scales. *Evol. Ecol. Res.*, 4, 993– 1004.
- Brannon, P. (2000). Niche relationships of two syntopic species of shrews, *Sorex fumeus* and *S. cinereus*, in the southern Appalachian Mountains. *Journal of Mammalogy*, 81(4):1053–1061.
- Brauer, F., and C. Castillo-Chavez (2000). *Mathematical Models in Population Biology and Epidemiology*, Springer-Verlag, Heidelberg.
- Breed, W. (1978). Ovulation rates and oestrus cycle lengths in several species of Australian rats (*Rattus* spp.) from various habitats. *Australian Journal of Zoology*, 26, 475-480.

- Breed, B. and F. Ford (2007). Native mice and rats. Natural History Series, CSIRO Publishing, Australia.
- Brown, J. (1973). Species diversity of seed-eating desert rodents in sand-dune habitats. *Ecology*, 54, 775-778.
- Brown, J. (1984). On the relationship between abundance and distribution of species. *Am. Nat.*, 124, 255–279.
- Brown, J., and B. Maurer. 1986. Body size, ecological dominance, and Cope's rule. *Nature* 324:248–250.
- Brown, J., and B. Maurer (1987). Evolution of species assemblages: effects of energetic constraints and species dynamics on the diversification of the American avifauna. *American Naturalist*, 130, 1-17.
- Brown, J., and G. Lieberman (1973). Resource Utilization and Coexistence of Seed-Eating Desert Rodents in Sand Dune Habitats *Ecology*, Vol. 54, No. 4, 788-797.
- Brown, J., Stevens, G., and D. Kaufman (1996). The geographic range: size, shape, boundaries, and internal structure. *Ann. Rev. Ecol. System.*, 27, 597–623.
- Bryce, J., Johnson, P., and D. Macdonald (2002). Can niche use in red and grey squirrels offer clues for their apparent coexistence? *Journal of Applied Ecology*, 39, 875-887.
- Bowers, M., and J. Brown (1982). Body Size and Coexistence in Desert Rodents: Chance or Community Structure? *Ecology*, Vol. 63, 391-400
- Buckland, S. (1980). A modified analysis of the Jolly-Seber capture-recapture model. *Biometrics* 36, 419-435.
- Burnett, S. (2008). Atherton Antechinus *Antechinus godmani*. In: S. Van Dyck and R. Strahan (eds). *The Mammals of Australia*. Reed New Holland, Australasia.
- Burnett, S., and Crowther (2008). (Rusty Antechinus *Antechinus adustus*. In: S. Van Dyck and R. Strahan (eds). *The Mammals of Australia*: Reed New Holland, Australasia.

- Bystrom, P., Andersson, J., Persson, L. and A. De Roos (2004). Size-dependent resource limitation and foraging-predation risk trade-offs: growth and habitat use in young arctic char. *Oikos* 104: 109–121.
- Buzas, M., Collins, L. and S. Culver. (2002) Latitudinal difference in biodiversity caused by higher tropical rate of increase. *Proceedings of the National Academy of Sciences* 99: 7841-7843.
- Calosi, Spicer, and Terblanche (2008). Phenotypic plasticity, dispersal, and physiology: determinants of range limits? [accessed 11 April 2009] <http://www2.plymouth.ac.uk/science/macrophysiology/abstracts/phenotypicplasticity.htm>.
- Campbell, N. (1996). Mitochondrial control region variation in two genera of Australian rodents; *Melomys* and *Uromys*: Application to phylogenetics, phylogeography and conservation. PhD Thesis, Centre for Conservation Technology, Southern Cross University.
- Campbell, N., and J. Reece, J.B (2008). *Biology*. 8<sup>th</sup> edition. Benjamin/Cummings Publishing Company, Inc.
- Cardillo, M. (2002a) Body size and latitudinal gradients in regional diversity of New World birds. *Global Ecology & Biogeography* 11 , 59–65.
- Cardillo, M. (2002b) The life-history basis of latitudinal diversity gradients: how do species traits vary from the poles to the equator? *Journal of Animal Ecology* 71, 79–87.
- Carter, R., and S. Prince (1988). Distributional limits from a demographic viewpoint. In: Davey, A., Hutchings, M., and Watkinson, A. (eds), *Plant Population Ecology*, Blackwell Scientific, Oxford, 165-184.
- Caughley, G., and A. Sinclair (1994). *Wildlife ecology and management*. Blackwell Science, Cambridge.
- Chao, A. (1987). Estimating the population size for capture-recapture data with unequal catchability. *Biometrics* 43(4), 783-792.

- Chao, A. and S. Lee (1992). Estimating the number of classes via sample coverage. *Journal of the American Statistical Association*, 87, 210-217.
- Chapman, M. (1999). Are there adequate data to assess how well theories of rarity apply to marine invertebrates. *Biodiversity and Conservation*, 8, 1295-1318.
- Chapple, D. (2003). Ecology, life-history, and behaviour in the Australian scincid genus *Egernia*, with comments on the evolution of complex sociality in lizards. *Herpetological Monographs*, 17, 2003, 145–180.
- Chase, J. (1996). Varying resources and competitive dynamics. *Am. Nat.* 147: 649-654.
- Chase, J., and M. Leibold (2003). *Ecological niches: Linking classical and contemporary approaches*. University of Chicago Press, Chicago.
- Chave, J., and E. Leigh (2002). A spatially explicit neutral model of biodiversity in tropical forests. *Theor. Popul. Biol.* 62, 153–168.
- Churchfield, S., and B. Sheftel (1994). Food niche overlap and ecological separation in a multi-species community of shrews in the Siberian taiga. *Journal of Zoology (London)*. 234(1), 105-124.
- Churchfield, S., Nesterenko, V., and E. Shvarts (1999). Food niche overlap and ecological separation amongst six species of coexisting forest shrews (Insectivora: Soricidae) in the Russian Far East. *Journal of Zoology*, 248: 349-359.
- Clarke, P., Kerrigan, R., and C. Westphal (2001). Dispersal potential and early growth in 14 tropical mangroves: do early life history traits correlate with patterns of adult distribution? *J. Ecol.*, 89, 648–659.
- Cody, M. (1986). Diversity, rarity, and conservation in Mediterranean-climate regions. In: M. Soule (ed). *Conservation Biology: the science of scarcity and diversity*. Sinauer Associates, Sunderland, MA., 122-152.

- Cofre, H., Böhning-Gaese, K., and P. Marquet<sup>1</sup> (2007a) Rarity in Chilean forest birds: which ecological and life-history traits matter? *Diversity and Distributions*, (*Diversity Distrib.*) 13, 203–212
- Cofre' Hernán L. Cofré, H., Samaniego, H., and P. Marquet (2007b). Patterns of Small Mammal Species Richness in Mediterranean and Temperate Chile. In: Kelt, Lessa, Salazar-Bravo, and Patton (Eds.). 275-302 *The Quintessential Naturalist: Honoring the Life and Legacy of Oliver P. Pearson*. University of California Publications in Zoology 134:1-981.
- Colwell, R. and E. Fuentes (1975). Experimental studies of the niche. *Annual Review of Ecology and Systematics* 6: 281-310.
- Colwell, R., Chang, K., and J. Chang (2004). Interpolating, extrapolating, and comparing incidence-based species accumulation curves. *Ecology*, 98, 2717-2727.
- Connell, J. (1978) Diversity in tropical rainforests and coral reefs. *Science* 199: 1302-1310.
- Connell, J. (1980). Diversity and the coevolution of competitors, or the ghost of competition past. *Oikos*, 35: 131–138.
- Cooper, W. and B. Cooper (2003). *Fruits of the rainforest: A guide to fruits in Australian tropical rainforest*. RD Press.
- Cormack, R. (1979). Models for capture-recapture. In: Cormack, R. M. & Patil, G. P. (eds). *Sampling biological populations*. International Cooperative Publishing House, Fairfield, Maryland.
- Cormack, R. (1989). Log-linear models for capture-recapture. *Biometrics* 45, 395-413.
- Cotgreave, P., and P. Harvey (1992). Relationships between body size, abundance and phylogeny in bird communities. *Functional Ecology*, 6, 248-256.
- Cotgreave, P., and M. Pagel (1997). Predicting and understanding rarity. In: (Eds) In: W. Kunin and K. Gaston (eds). *The Biology of rarity, Population and Community Biology Series*, Chapman and Hall, London, 262-272.



- Cottenie, K., and L. De Meester (2004) Metacommunity structure: synergy of biotic interactions as selective agents and dispersal as fuel. *Ecology* 85:114–119.
- Coyne, J., and H. Orr (2004). *Speciation*. Sinauer Associates, Inc. Sunderland, Massachusetts, USA.
- Craig Roberts, S. (2007). Scent marking. In: *Rodent societies: An ecological and Evolutionary Perspective* (Eds) Wolff and Sherman, University of Chicago Press.
- Crawley, M. (1992). Seed predators and plant population dynamics. In: M. Fenner (Ed), *Seeds: the ecology of regeneration in plant communities*, 157–191. CAB International, Wallingford, UK.
- Crawley, M. (1997). Plant–herbivore dynamics. In: *Plant ecology*. Second edition (Ed) Crawley, 401– 474 Blackwell Scientific Publications, Oxford, UK.
- Crawley, M. (2000). Seed predators and plant population dynamics. In: *Seeds: the ecology of regeneration in plant communities: Second edition*. (Ed) Fenner, 167–182. CAB International, Wallingford, UK.
- Crome, F., Isaacs, J., and L. Moore (1994). The utility to birds and mammals of remnant riparian vegetation and associated windbreaks in the tropical Queensland uplands. *Pacific Conservation Biology*, 1, 328-343.
- Crome, F., Moore, L., and G. Richards (1991). A study of logging damage in upland rainforest in north Queensland. *Forest Ecology and Management*, 49, 1-29.
- Crome, F., Thomas, M., and L. Moore (1994). A novel Bayesian way to assessing impacts of rainforest logging. Commonwealth Scientific and Industrial Research Organisation (CSIRO) Memorandum.
- Crowson, R. (1981). *The biology of the Coleoptera*. Academic Press, London.
- Crowther, M. (2008). Yellow-footed Antechinus *Antechinus flavipes*. In: S. Van Dyck and R. Strahan (eds). *The Mammals of Australia*, Reed New Holland, Australasia.

- Damuth, J. (1981). Population density and body size in mammals. *Nature*, 290, 699–700.
- Damuth, J. (1991). Of size and abundance. *Nature*, 351, 268–269.
- Damuth, J. (1993). Cope's rule, the island rule, and the scaling of mammalian population density. *Nature*, 365, 748–750.
- Darroch, J. (1958). The multiple recapture census: Estimation of a closed population. *Biometrika* 45, 343-351.
- Darwin, C. (1859), *On the Origin of Species by Means of Natural Selection, or the Preservation of Favoured Races in the Struggle for Life*, London: John Murray.
- Davies, K., Margules, C., and J. Lawrence (2004). A synergistic effect puts rare, specialized species at greater risk of extinction. *Ecology*, Vol. 85, No. 1, 265-271.
- Dennis A. and P. Johnson (2008). Musky Rat-Kangaroo *Hypsiprymnodon moschatus*. In: S. Van Dyck and R. Strahan (eds). *The Mammals of Australia*, Reed New Holland, Australasia.
- Dennis, A. (1997). Musky Rat-kangaroos, *Hypsiprymnodon moschatus*, cursorial frugivores in Australia's wet-tropical rain-forests. Ph.D. Thesis, James Cook University of North Queensland, Townsville, Australia.
- Diamond, J. (1975). Assembly of species communities. In: (eds) Cody, M. and Diamond, J., *Ecology and evolution of communities*. Belknap/Harvard Univ. Press, 342-444.
- Dice, L. (1938). Some census methods for mammals. *J. Wildl. Manage.* 2, 119-130.
- Dice, L. (1941). Methods for estimating populations of animals. *J. Wildl. Manage.* 5, 398-407.
- Dickman, C. and E. Stodart (2008). Long-nosed Bandicoot *Perameles nasuta*. In: S. Van Dyck and R. Strahan (eds). *The Mammals of Australia*, Reed New Holland, Australasia.

- Dobson, F., and J. Yu (1993). Rarity in Neotropical Forest Mammals Revisited. Conservation Biology Volume 7, No. 3.
- Dodge, W. (1967). Biology and life history of the porcupine (*Erethizon dorsatum*) in western Massachusetts. PhD. thesis. Univ. of Mass., Amherst, Mass.
- Dodge, W. (1982). Porcupine. In "Wild Mammals of North America: Biology, Management and Economics." Chapman, J.A.; Feldhamer, G.A., eds. Johns Hopkins University Press, Baltimore, MD.
- Doherty, D., and A. Harcourt (2004). Are rare primate taxa specialists or simply less studied? Journal of Biogeography, (31), 57 – 61.
- Donnellan, S. (1989). Phylogenetic relationships of New Guinea rodents (Rodentia: Muridae) based on chromosomes. Australian Mammalogy, 12, 61-67.
- Drury, W. (1974). Rare species. Biological Conservation, 6, 162-169.
- Dunham, J., and G. Vinyard (1997) Relationships between body mass, population density, and the self-thinning rule in stream-living salmonids. Can. J. Fish. Aquat. Sci. 54: 1025-1030.
- Duralia, T., and R. Reader (1993). Does abundance reflect competitive ability? A field test with three prairie grasses. Oikos, 68, 82-90.
- Edwards, G., and M. Crawley (1999). Rodent seed predation and seedling recruitment in mesic grassland. Oecologia 118: 288-296.
- Edwards, J., Heckel, D., and D. Guynn (1985). Niche overlap in sympatric populations of fox and grey squirrels. Journal of Wildlife Management, 62, Vol. 1, 354-363.
- Edwards, W., and M. Westoby (1996). Reserve mass and dispersal investment in relation to geographic range of plant species: phylogenetically independent contrasts. J. Biogeogr., 23, 329–338.
- Efford, M. (2004). Density estimation in live-trapping studies. OIKOS 106: 598-610.
- Ernest, S. (2005). Body Size, Energy Use, and Community Structure of Small Mammals. Ecology, Vol. 86, No. 6, 1407-1413.

- Estes, J., Jameson, R., and E. Rhode (1982). Activity and prey selection in the sea otter: influence of population status on community structure. *American Naturalist*, 120, 242-258.
- Etienne, R., Alonso, D., and J. McKane (2001). The zero-sum assumption in neutral biodiversity theory. *Journal of Theoretical Biology* 248, 522–536
- Etienne, R., and H. Olff (2004). A novel genealogical approach to neutral biodiversity theory. *Ecol. Lett.*, 7, 170–175.
- Fiedler, P. (1987). Life History and Population Dynamics of Rare and Common Mariposa Lilies (*Calochortus* Pursh: Liliaceae). *Journal of Ecology*, Vol. 75, No. 4, 977-995.
- Fisher, R., Corbet, A., and C. Williams (1943). The relation between the number of species and the number of individuals in a random sample of an animal population. *Journal of Animal Ecology*, 12, 42-58.
- Flannery, T. (1995). *Mammals of the South-West Pacific and Moluccan Islands*. Reed Books, Sydney.
- Flannery, T., and P. Schouten (2001). *A Gap in Nature: Discovering the World's Extinct Animals*. Atlantic Monthly Press, New York.
- Ford, F. (2003). Conilurine rodent evolution: The role of ecology in modifying evolutionary consequences of environmental change. PhD Thesis, School of Zoology and Tropical Ecology, James Cook University, Townsville, Queensland.
- Ford, H. (1990). Relationships between distribution, abundance, and foraging specialisation in Australian landbirds. *Ornis Scandinavica*, 21, 133-138.
- Frankham, R. (1996). Relationship of genetic variation to population size in wildlife. *Conserv. Biol.* 10, 1500-1508.
- Frankham, R. (1997). Do island populations have lower genetic variation than mainland populations? *Heredity* 78, 311-327.

- Frankham, R., Ballou, J., and D. Briscoe (2002). *Introduction to conservation genetics*. Cambridge University Press, UK.
- Frankham, R., Gilligan, D., Morris, D., and D. Briscoe (2001). Inbreeding and extinction: effects of purging. *Conserv. Genet.* 2, 279-284.
- Freeland, W. (1972). *A rainforest and its rodents*. Masters of Science thesis, Ecology Department, University of Queensland.
- Fretwell, S., and Lucas (1970). On territorial behaviour and other factors influencing habitat distribution of birds. *Theoretical developments. Acta Biotheoretica*, 19, 16-36.
- Gaston, K. (1991). How big is a species' geographic range? *Oikos*, 61, 434-438.
- Gaston, K. (1994). *Rarity. Population and Community Biology Series*, Chapman and Hall, London.
- Gaston, K. (1996). Species-range-size distributions: patterns, mechanisms and implications. *Trends Ecol. Evol.*, 11, 197-201.
- Gaston, K. (2003). *The Structure and Dynamics of Geographic Ranges*. Oxford University Press, Oxford.
- Gaston, K., and S. Chown (2005). Neutrality and the niche. *Functional Ecology* 19, 1-6.
- Gaston, K., and J. Lawton (1988). Patterns in body size, population dynamics, and regional distribution of bracken herbivores. *American Naturalist*, 132, 662-680.
- Gaston, K., and R. Fuller (2008). Commonness, population depletion and conservation biology. *Trends in Ecology & Evolution* Volume 23, (1), 14-19.
- Gaston, K., and T. Blackburn (1996). Some conservation implications of geographic range size-body size relationships. *Conservation Biology*, 10, 638-646.
- Gaston, K., and T. Blackburn (2000). *Pattern and Process in Macroecology*. Blackwell Science, Malden, MA.

- Gaston, K., and W. Kunin (1997). Concluding comments. In: W. Kunin and K. Gaston (eds). *The Biology of rarity, Population and Community Biology Series*, Chapman and Hall, London, 262-272.
- Gause, G. (1934). *The struggle for existence*. Williams and Wilkins, Baltimore.
- Genner, M., Turner, G., and S. Hawkins (1999). Foraging of rocky habitat cichlid fishes in Lake Malawi: coexistence through niche partitioning? *Oecologia* 121 (2): 283-292.
- Gilbert, B., and M. Lechowicz. (2004). Neutrality, niches, and dispersal in a temperate forest understory. *Proceedings of the National Academy of Sciences, USA* 101:7651–7656.
- Gilliam, J, and D. Fraser (1987). Habitat selection under predator hazard: test of a model with foraging minnows. *Ecology* 68: 1856-1862.
- Glanz, W. (1984). Ecological relationships of 2 species of *Akodon* in central Chile. *Journal of Mammalogy*. 65(3), 433-441.
- Glazier, D. (1980). Ecological shifts and the evolution of geographically restricted species of North American *Peromyscus* (mice). *Journal of biogeography*, 7, 63-83.
- Glazier, D. (1986). Temporal variation of abundance and the distribution of species. *Oikos*, 47, 309-314.
- Goodwin, N., Dulvy, N., and J. Reynolds (2005). Macroecology of live-bearing in fishes: latitudinal and depth range comparisons with egg-laying relatives. *Oikos*, 110, 209–218.
- Goosem, M. (2000). Effects of tropical rainforest roads on small mammals: edge changes in community composition *Wildlife Research*, 2000, 27, 151–163.
- Goosem, M. (2002) Effects of tropical rainforest roads on small mammals: fragmentation, edge effects and traffic disturbance. *Wildlife Research*, 29, 277 – 289.

- Gordon, G. (2008). Northern Brown Bandicoot *Isoodon macrourus*. In: S. Van Dyck and R. Strahan (eds). The Mammals of Australia. Reed New Holland, Australasia.
- Grand, T. (2002). Alternative forms of competition and predation dramatically affect habitat selection under foraging–predation-risk trade-offs. *Behavioural Ecology* Vol. 13 No. 2: 280–290.
- Green, M. (1987). Ecological separation in Himalayan ungulates. *Journal of Zoology Series B*. 1(4), 693-720.
- Green, R. (1979). The ecology of wood mice (*Apodemus sylvaticus*) on arable farmland. *J. Zool.*, 188:357-377.
- Griesemer, J. (1992). Niche: Historical perspectives. In: *Keywords in evolutionary biology*. (Eds) Fox-Keller and Lyoyd, 231-240, Cambridge: Harvard University Press.
- Griggs, R. (1940). The ecology of rare plants. *Bulletin of the Torrey Botanical Club* 67, 575-594.
- Groeters, F. and D. Shaw, D. (1992) Association between latitudinal variation for embryonic development time and chromosome structure in the grasshopper *Caledia captiva* (Orthoptera: Acrididae). *Evolution* 46:245–257.
- Groves, C., and T., Flannery (1994). A revision of the genus *Uromys* Peters, 1867 (Muridae: Mammalia) with descriptions of two new species. *Records of the Australian Museum*, Vol. 46, 145-169.
- Gutierrez, D., and R. Menendez (1997). Patterns in the distribution, abundance and body size of carabid beetles (Coleoptera: Caraboidea) in relation to dispersal ability. *J. Biogeogr.*, 24, 903– 914.
- Haffenden, A., and R. Atherton (2008). Long-tailed Pygmy Possum *Cercartetus caudatus*. In: S. Van Dyck and R. Strahan (eds). *Mammals of Australia*, Reed New Holland, Australasia.

- Halpin, Z. (1987). Natal dispersal and formation of new social groups in a newly established town of black-tailed prairie dogs (*Cynomys ludovicianus*). In: B. Chepko-Sade and Z. Halpin (eds), *Mammalian dispersal patterns: The effects of social structure on population genetics*, University of Chicago Press, 104-118.
- Hansen, T. (1980). Influence of larval dispersal and geographic distribution on species longevity in neogastropods. *Paleobiology*, 6, 193–207.
- Hanski, I. (1982). Dynamics of regional distribution: the core-satellite species hypothesis. *Oikos*, 38, 210-221.
- Hanski, I. (1985). Single-Species Spatial Dynamics May Contribute to Long-Term Rarity and Commonness Ecology, Vol. 66, No. 2, 335-343.
- Hanski, I. (1991), Single-species metapopulation dynamics: concepts, models, and observations. *Biological Journal of the Linnean Society*, 42, 17-38.
- Hanski, I. (1999). *Metapopulation Ecology*. Oxford University Press, New York.
- Hanski, I., Kouki, J. and A. Halkka. (1993). Three explanations of the positive relationship between distribution and abundance of species. In: Ricklefs and Schluter (eds) *Species Diversity in Ecological Communities: Historical and Geographical Perspectives*. University of Chicago Press, Chicago, 108–116.
- Hanski, I., Kouki, J., and A. Halkka (1993). Three explanations of the positive relationship between distribution and abundance of species. In: Ricklefs, R., and Schluter, D. (eds). *Historical and Geographical determinants of Community Diversity*. University of Chicago Press, Chicago.
- Hansen, T. (1980). Influence of Larval Dispersal and Geographic Distribution on Species Longevity in Neogastropods. *Paleobiology*, Vol. 6, 193-207.
- Hanson, A. (2006). History of the Norway rat (*Rattus norvegicus*): Rat behaviour and biology. [accessed 21 August 2007]. <http://www.ratbehavior.org/history.htm>.
- Hansson, L. (1987). An interpretation of rodent dynamics as due to trophic interactions. *Oikos* 50: 308-318.



- Harcourt, A. (2006). Rarity in the tropics: biogeography and macroecology of the primates. *Journal of Biogeography*, 33, 2077–2087.
- Harcourt, A., S. Coppeto and S. Parks (2002). Rarity, specialization and extinction in primates. Volume 29 (4), 445 – 456.
- Harder, L. (1979). Winter feeding by porcupines in montane forests of southwestern Alberta. *Can. Field Nat.* 93:405-410.
- Hardin, G. (1960). The Competitive Exclusion Principle. *Science* 131, 1292-1297
- Harper, J. (1981). The meanings of rarity. In: *The biological aspects of rare plant conservation* (ed) H. Synge, Wiley, New York, 189-203.
- Harrington, G., Irvine, A., Crome, F., and L. Moore (1997). Large-seeded trees in rainforest fragments: a study of higher order interactions. pp.292-303 in Laurance, W. F., and Bierregaard R. O. (eds). *Tropical forest remnants: ecology, management and conservation of fragmented assemblages*. University of Chicago Press, Chicago.
- Harrington, G., Freeman, A., and F. Crome (2001). The effects of fragmentation of an Australian tropical rain forest on populations and assemblages of small mammals *Journal of Tropical Ecology* 17:225-240
- Harvey, P., and M. Pagel (1991). *The comparative method in evolutionary biology*. Oxford University Press, Oxford.
- Hedderson, T. (1985). Rarity at its range limits: dispersal capacity and habitat relationships of extraneous moss species in a boreal Canadian National Park. *Biological Conservation* 59, 113-120.
- Hegazy, A. and N. Eesa (1991). On the Ecology, Insect Seed-Predation, and Conservation of a Rare and Endemic Plant Species: *Ebenus armitagei* (Leguminosae). *Conservation Biology*, Vol. 5, 317-324.
- Heinsohn, G., and R. Heinsohn (1999). Long-term dynamics of a rodent community in an Australian tropical rainforest. *Wildlife Research*, 1999, 26, 187-198.

- Heinsohn, T., and G. Hope (2006). The Torresian connections: Zoogeography of New Guinea. In: Merrick, J., Archer, M., Hickey, G., and Lee, M. (eds). Evolution and biogeography of Australasian vertebrates. Auscipub Pty Ltd. NSW.
- Hilbert D., Ostendorf B., and M. Hopkins (2001) Sensitivity of tropical forests to climate change in the humid tropics of north Queensland. *Austral Ecology*, 26, 590–603.
- Holt, R. (1997). Rarity and evolution: some theoretical considerations. In: *The biology of rarity*. Edited by William Kunin and Kevin Gaston. Chapman and Hall, London.
- Holt, R., and T. Keitt (2000). Alternative causes for range limits: a metapopulation perspective. *Ecol. Lett.*, 3, 41–47.
- Hope, G. (1996). Quaternary change and historical biogeography of Pacific Islands. Pp. 165-190. In: Keast, A., Miller, S. (eds). *The origin and Evolution of Pacific Island Biotas, New Guinea to Eastern Polynesia: Patterns and Processes*. SPB Academic Publishing, Amsterdam.
- Hopkins, W. (2001) *A New View of Statistics*: [accessed 23 March 2009] [http://davidmlane.com/hyperstat/logic\\_hypothesis.html](http://davidmlane.com/hyperstat/logic_hypothesis.html).
- Houston, A., McNamara, J., and J. Hutchinson (1993). General Results concerning the Trade-Off between Gaining Energy and Avoiding Predation. *Philosophical Transactions: Biological Sciences*, Vol. 341, 375-397.
- Hubbell, S. (1997). A unified theory of biogeography and relative species abundance and its application to tropical rain forests and coral reefs, *Coral Reefs* 16, S9–S21.
- Hubbell, S.P. (2001). *The Unified Neutral Theory of Biodiversity and Biogeography*. Princeton University Press.
- Hubbell, S. and J. Lake. (2003). The neutral theory of biodiversity and biogeography, and beyond. In: *Macroecology: Patterns and Processes* (eds) Blackburn, T. & Gaston, K.). Blackwell, Oxford.

- Hubbell, S. (2006). Neutral theory and the evolution of ecological equivalence. *Ecology* 87:1387–1398.
- Huitu, O. (2003) Temporal and spatial synchrony in small mammal population fluctuations. Phd Thesis, Turun Yliopiston Julkaisuja, Annales Universitatis Turkuensis
- Hurlbert, S. (1971) The non-concept of species diversity: a critique and alternative parameters. *Ecology* 52, 577-586.
- Hutchinson, G. (1957). Concluding remarks. *Cold Spring Harbor Symposium Quantitative Biology* 22: 415-427.
- Hutchinson, G. (1959). Homage to Santa Rosalia, or why are there so many kinds of animals? *American Naturalist*, 93, 145-59.
- Hutchinson, G. (1978). *An introduction to population ecology*. New Haven: Yale University Press.
- Hyland, B., Whiffin, T., Christophel, D., Gray, B., and E. Elick. (2003). *Key to Australian tropical rainforest plants*. CSIRO Publishing, Melbourne, Australia.
- J´acomo, Silveira, L., and J. Felizola (2004). Niche separation between the maned wolf (*Chrysocyon brachyurus*), the crab-eating fox (*Dusicyon thous*) and the hoary fox (*Dusicyon vetulus*) in central Brazil. *J. Zool., Lond.* 262, 99–106.
- Jablonski, D. (1986). Larval ecology and macroevolution in marine invertebrates. *Bull. Mar. Sci.*, 39, 565–587.
- Jablonski, D. (1987). Heritability at the species level analysis of geographic ranges of Cretaceous mollusks. *Science*, 238, 360– 363.
- Jablonski, D. and K. Roy (2003). Geographical range and speciation in fossil and living molluscs. *Proc. Roy. Soc. London Ser. B*, 270, 401–406.
- Jablonski, D., Flessa, K., and J. Valentine (1985). Biogeography and paleobiology. *Paleobiology*, 11, 75–90.
- Jett, D., and J. Nichols (1987). A field comparison of nested grid and trapping web density estimators. *J. Mammal.* 68, 888-892.

- Johnson, C. (1998). Rarity in the tropics: latitudinal gradients in distribution and abundance in Australian mammals. *Journ. Anim. Ecol.*, 67, 689-698.
- Johnson, C., and K. Vernes (2008). Red-legged Pademelon *Thylogale stigmatica*. In: S. Van Dyck and R. Strahan (eds). *The Mammals of Australia*. Reed New Holland, Australasia.
- Johnson, M., and P. Raven (1973). Species number and endemism: The Galapagos Archipelago revisited, *Science*, 179, 893-895
- Jolly, G. (1982). Mark-recapture models with parameters constant in time. *Biometrics* 38, 301-321.
- Jones, M., and L. Barmuta (2000). Niche differentiation among sympatric Australian dasyurid carnivores. *Journal of Mammalogy*, 81(2):434–447.
- Juliano, S. (1983). Body size, dispersal ability, and range size in North American species of *Brachinus* (Coleoptera: Carabidae). *Coleopt. Bull.*, 37, 232–238.
- Karanth, K., and J. Nichols (1998). Estimation of tiger densities in India using photographic captures and recaptures. *Ecology* 79, 2852-2862.
- Karron, J. (1987). The pollination ecology of co-occurring geographically restricted and widespread species of *Astragalus* (Fabaceae). *Biological Conservation*, 39, 179-193.
- Kattan, G. (1992). The birds of the Cordillera Central of Colombia. *Conservation Biology* 6, 64-70.
- Kaufman, D. (1995) Diversity of New World mammals: universality of the latitudinal gradients of species and bauplans. *Journal of Mammalogy* 76: 322-334.
- Kavanaugh, D. (1985). On wing atrophy in carabid beetles (Insecta: Coleoptera: Carabidea), with special reference to Nearctic *Nebria*. In: *Taxonomy, Phylogent, and zoogeography of beetles and ants* (Ed) Ball. Dr W. Junk Publ., Dordrecht.
- Kawano, K. (2002). Character displacement in giant rhinoceros beetles. *Am. Nat.* 159, 255-271.

- Keast, A., (1961). Bird speciation on the Australian continent. *Bulletin of the museum of Comparative Zoology*, 123, 303-495.
- Keast, A. (1968). Competitive Interactions and the Evolution of Ecological Niches as Illustrated by the Australian Honeyeater Genus *Melithreptus* (Meliphagidae). *Evolution*, Vol. 22, No. 4, 762-784.
- Kerle, J. (2008). Grassland Melomys *Melomys burtoni*. In: S. Van Dyck and R. Strahan (eds). *Mammals of Australia*, Reed New Holland, Australasia.
- Kincaid, B., Cameron, G., and A. Carnes (1983). Patterns of Habitat Utilization in Sympatric Rodents on the Texas Coastal Prairie: *Ecology*, 64, 1471-1480.
- Kiltie, R. (1982). Bite force as a basis for niche differentiation between rain forest Peccaries (*Tayassu tajacu* and *T. pecari*). *Biotropica*, Vol. 14, No. 3.
- Kinlan, B., and A. Hastings (2005). Rates of population spread and geographic range expansion: what exotic species tell us. In: *Species Invasions: Insights to Ecology, Evolution and Biogeography* (eds Sax, D., Gaines, S.D. & Stachowicz, J.J.). Sinauer Associates, Sunderland, MA, 381–419.
- Knox, A., Losos, B., and C. Schneider (2001). Adaptive radiation versus intraspecific differentiation: morphological variation in Caribbean Anolis lizards. *J. Evol. Biol.* 14: 904-909.
- Kouki, J., and U. Hūyrynen (1991). On the relationship between distribution and abundance in birds breeding on Finnish mires: the effect of habitat specialisation. *Ornis fennica*, 68, 170-177.
- Kozakiewicz, M., and A. Kozakiewicz (2004). Chemical information, scent trails and spatial behaviour of small forest rodents. a review, *Ejpa* 7(2), #03.
- Krebs, C. (2005). *Ecological Methodology*. Harper Collins, New York.
- Kruckenber, A., and D. Rabinowitz (1985). Biological aspects of endemism in higher plants. *Annual review of Ecology and Systematics*, 16, 447-479.
- Kryštufek, B., and H. Griffiths (2002). Species richness and rarity in European rodents. *Ecography*, Vol. 25, Issue 1, 120 – 128.

- Kunin, W. (1997). Introduction: on the causes and consequences of rare-common differences. In: W. Kunin and K. Gaston (eds). *The Biology of rarity, Population and Community Biology Series*, Chapman and Hall, London.
- Kunin, W., and K. Gaston (1993). The biology of rarity: patterns, causes, and consequences. *Trends in Ecology and Evolution*, 8, 298-201.
- Lack, D. (1957). *Darwin's finches*. Cambridge University Press.
- Lahti, T., Kempainen, E., Kurtto, A., and P. Uotila (1991). Distributional and biological characteristics of threatened vascular plants in Finland. *Biological Conservation*, 55, 299-314.
- Larsen, D. (1998). Mark and recapture methods. [Accessed 10 June 2007] <http://people.umass.edu/dsigourn/jolly-Seber%20method.pdf>.
- Laurance, W. (1989). Ecological impacts of tropical forest fragmentation on non-flying mammals and their habitats. PhD Thesis. University of California, Berkely.
- Laurance, W. F. (1991). Ecological correlates of extinction proneness in Australian tropical rainforest mammals. *Conservation Biology* 5, 79-89.
- Laurance, W. (1992). Abundance Estimates of Small Mammals in Australian Tropical Rainforest: a Comparison of Four Trapping Methods *Wildl. Res.*, 19, 651-655.
- Laurance, W., Garesche, J., and C. Payne (1993). Avian nest predation in modified and natural habitats in tropical Queensland: An experimental study. *Wildl. Res.*, 20, 711 -723.
- Lawler, S., and P. Morin (1993). Food web architecture and population dynamics in laboratory microcosms of protists. *Am. Nat.* 141: 675-686.
- Lawton, J. (1989). What is the relationship between population density and body size in animals? *Oikos*, 55, 429-434.
- Leibold, M. (1995). The niche concept revisited: Mechanistic models and community context. *Ecology* 76:1371-1382.

- Leslie, P. (1952). The estimation of population parameters from data obtained by means of the capture-recapture method II. The estimation of total numbers. *Biometrika* 39.
- Leslie, P., Chitty, D., and H. Chitty (1953). The estimation of population parameters from data obtained by means of the capture recapture method , III: an example of the practical applications of the method. *Biometrika* 40: 137-169.
- Lester, S., and B. Ruttenberg (2005). The relationship between pelagic larval duration and range size in tropical reef fishes: a synthetic analysis. *Proc. Roy. Soc. London Ser. B*, 272, 585–591.
- Lester, S., Ruttenberg, B., Gaines, S., and B. Kinlan (2007). The relationship between dispersal ability and geographic range size. *Ecology Letters* 10: 745–758.
- Leung, L. (1994). Ecology of three species of small mammals in the iron Range area, Cape York Peninsula: what factors influence the populations? PhD Thesis, School of Biological Sciences, University of Sydney, NSW, Australia.
- Leung, L., Dickman, C., and L. Moore (1993). Genetic variation in fragmented populations of a rainforest rodent, *Melomys cervinipes*. *Pacific Conservation Biology*, (1), 58-65.
- Levine, J., and J. HilleRisLambers (2009). The importance of niches for the maintenance of species diversity. *Nature*, 2009.
- Levins, R. (1969). Some demographic and genetic consequences of environmental heterogeneity for biological control. *Bull. Entomol. Soc. Am.*, 15, 237–240.
- Levinton, J. (1983). The latitudinal compensation hypothesis: growth data and a model of latitudinal growth differentiation based upon energy budgets: Interspecific comparison based on *Ophryotrocha puerlis* (Polychaeta: Dorvilleidae). *Biol Bull.* 165:686–698.
- Likert, R. (1932). A Technique for the Measurement of Attitudes, *Archives of Psychology*, No.140.

- Lima, S., and L. Dill (1990). Behavioral decisions made under the risk of predation: a review and prospectus. *Can. J. Zool.* 68(4): 619–640.
- Lister, B. (1976). The nature of niche expansion in West Indian *Anolis* lizards: Ecological consequences of reduced competition. *Evolution* 30: 659-676.
- Lloyd, K., Wilson, J., and W. Lee (2003). Correlates of geographic range size in New Zealand *Chionochloa* (Poaceae) species. *J. Biogeogr.*, 30, 1751–1761.
- Loeuille, N., and Loreau, M. (2006). Evolution of body size in food webs: does the energetic equivalence rule hold? *Ecology Letters*, 9: 171–178.
- Lotka, A. (1932). The growth of mixed populations: Two species competing for the common food supply. *Journal of the Washington Academy of Sciences*. 21, 461-469.
- Louette, M., and P. Herroelen (2007). Comparative biology of the forest-inhabiting hawks *Accipiter* spp. in the Democratic Republic of Congo. *Ostrich*. 78(1). 21-29.
- Lowry, E., and S. Lester (2006). The biogeography of plant reproduction: potential determinants of species range sizes. *J. Biogeogr.*, 33, 1975–1982.
- Lubchenco, J., and B. Menge (1978) Community Development and Persistence in a Low Rocky Intertidal Zone. *Ecological Monographs*: Vol. 48, No. 1, 67-94.
- Lunney, D. (2008). Bush Rat *Rattus fuscipes*. In: S. Van Dyck and R. Strahan (eds). *The Mammals of Australia*: Reed New Holland, Australasia.
- MacArthur, R., and R. Levins (1964). Competition, habitat selection, and character displacement in a patchy environment. *Proc. Nat. Acad. Sci.* 51, 1207-1210.
- MacArthur, R., and E. Wilson (1967). *The theory of island biogeography*. Princeton,: Princeton Press.
- MacArthur, R. (1955). Fluctuations of animal populations and a measure of community stability. *Ecology* 36: 533-536.
- MacArthur, R. (1969). The theory of the niche. In: (Eds) *Population biology and evolution*. 159-176. Syracuse University Press.



- MacArthur, R. (1972). *Geographical Ecology*. Harper and Row, New York.
- McCauley, S., Davis, C., Relyea, R., Yurewicz, K., Skelly, K., and E. Werner (2008).  
Metacommunity patterns in larval odonates. *Oecologia* (2008) 158:329–342
- McGill, B. (2003) A test of the unified neutral theory of biodiversity. *Nature* 422, 881–885.
- Macgowan, J., and P. Walker (1979). Structure in the copepod community of the North Pacific Central Gyre. *Ecological Monographs*, 49, 195-226.
- Marchlewska-Koj A. (2000). Olfactory and ultrasonic communication in bank voles. *Pol. J. Ecol.*, 48 (Suppl.), 11-20.
- McKane, A., Alonso, D., and R. Sole´ (2004). Analytic solution of Hubbell’s model of local community dynamics. *Theor. Popul. Biol.*, 65, 67–73.
- MacNally, R. (1989). The relationship between habitat breadth, habitat position, and abundance in forest and woodland birds along a continental gradient. *Oikos* 54: 44-54.
- MacNally, R. and J. Doolan (1986). An empirical approach to guild structure: habitat relationships in nine species of eastern-Australian cicadas. *Oikos* 47: 33-46.
- MacNally, R., and G. Brown (2001). Reptiles and habitat fragmentation in the Box-ironbush Forests of Central Victoria, Australia: Predicting Compositional Change and Faunal Nested-ness. *Oecologia* 128:116-125.
- Maehr, D. (1997). The comparative ecology of bobcat, black bear, and Florida panther in South Florida. *Bulletin of the Florida Museum of Natural History*. 40(1). 1-176.
- Malmqvist, B. (2000). How does wing length relate to distribution patterns of stoneflies (Plecoptera) and mayflies (Ephemeroptera)? *Biol. Conserv.*, 93, 271–276.
- Margules, C., and Usher, M. (1981). Criteria used in assessing wildlife conservation potential: a review. *Biological Conservation*, 21, 79-109.
- Margules, D., and C., Lawrence (2004). A synergistic effect puts rare specialised species at greater risk of extinction. *Ecology*, 85 (1), 265-271.

- Maurer, B., Ford, J., and Rapoport, E. (1991). Extinction rate, body size, and avifaunal diversity. *Acta XX Congressus Internationalis Ornithologici*, 826-834.
- May, R. (1975). Patterns of species abundance and diversity. In: M.L. Cody and J. M. Diamond, (eds)., *Ecology and evolution of communities*. Harvard University press.
- Mayr, E. (1963). *Animal species and evolution*. The Belknap Pres of Harvard University Press, Cambridge, MA.
- McAllister, D., Platania, S., Schueler, F. (1986). Ichthyofaunal patterns on a geographical grid,. In: C. Hocutt and E. Wiley (eds). *Zoogeography in Freshwater Fishes of North America*. Wiley, New York, 17-51.
- McCoy, E., and Mushinsky, H. (1992). Rarity of organisms in the sand pine-scrub habitat of Florida. *Conservation Biology*, 6, 537-548.
- McDonald, L. (2004). Sampling rare populations. In: *Sampling Rare and Elusive Species* (Ed) William L. Thompson. Island Press, London.
- M'Closkey, R. (1976). Community Structure in Sympatric Rodents. *Ecology*, Vol. 57, No. 4), 728-739.
- M'Closkey, R., and Fieldwick, B. (1975). Ecological Separation of Sympatric Rodents (*Peromyscus* and *Microtus*). *Journal of Mammalogy*, Vol. 56, No. 1. 119-129.
- Miller, R. (1967). Patterns and process in competition. *Adv. Ecol. Res.* 4: 1-74.
- Monro, J. (1967). The Exploitation and Conservation of Resources by Populations of Insects. *Journal of Animal Ecology*, Vol. 36, 531-547.
- Moore, L. (2008). Giant White-tailed Rat *Uromys caudimaculatus*. In: In: S. Van Dyck and R. Strahan (eds). *The Mammals of Australia*, Reed New Holland, Australasia.
- Moore, L., and S. Burnett (2008). Fawn-footed Melomys *Melomys cervinipes*. In: In: S. Van Dyck and R. Strahan (eds). *The Mammals of Australia*, Reed New Holland, Australasia.
- Moore, L., and L. Leung (2008). Cape York Rat *Rattus leucopus*. In: S. Van Dyck and

- R. Strahan (eds). The Mammals of Australia. Reed New Holland, Australasia.
- Moore, L., and J. Winter (2008). Pygmy White-tailed Rat *Uromys hadrourus*. In: S. Van Dyck and R. Strahan (eds). The Mammals of Australia, Reed New Holland, Australasia.
- Moore, N., and D. Gillieson (2008). Using field survey and remote sensing to assess rainforest canopy damage following Cyclone Larry. *Austral Ecology* 33, 417–431.
- Mora, C., and D. Robertson (2005). Factors shaping the range-size frequency distribution of the endemic fish fauna of the tropical eastern Pacific. *J. Biogeogr.*, 32, 277–286.
- Morcombe, M. (2000). Field guide to Australian birds. Steve Parish Publishing.
- Morse, D., Stork, N., and J. Lawton. (1988). Species number, species abundance, and body length relationships of arboreal beetles in Bornean lowland rainforest trees. *Ecological Entomology*, 13, 25-37.
- Müller-Schwarze, D. and S. Heckman (1980). The social role of scent marks in beaver. *Castor canadensis*. *Journal of Chemical Ecology*, 6:81-95.
- Müller-Schwarze, D. (1992). Castoreum of beaver (*Castor canadensis*): Function, chemistry, and biological activity of its components. In: *Chemical signals in vertebrates 6* (Eds) Doty and Müller-Schwarze: 457-464, New York: Plenum.
- Nee, S., Read, A., Greenwood, J., and P. Harvey (1991). The relationship between abundance and body size in British birds. *Nature* 351, 312-313.
- Oakwood, M., Jurado, E., Leishman, M., and M. Westoby (1993). Geographic ranges of plant species in relation to dispersal morphology, growth form and diaspore weight. *J. Biogeogr.*, 20, 563–572.
- Odum, E., and H. Odum. (1959). *Fundamentals of Ecology*. W.B. Saunders, Philadelphia.

- Ooi, M. (2007). Comparative ecology of rare and common species in a fire-prone system. PhD Thesis, University of Wollongong University of Wollongong, Thesis Collections.
- Orians, G. (1997). Evolved consequences of rarity. In: W. Kunin and K. Gaston (eds). *The Biology of rarity*, Population and Community Biology Series, Chapman and Hall, London.
- Otis, D., Burnham, K., White, G., and D. Anderson (1978). Statistical inference from capture data on closed animal populations. *Wildlife Monographs* 62: 1-135.
- Paine, M. (1990). Life history of darters (Percidae: Etheostomatiini) and their relationship with body size, reproductive behaviour, latitude, and rarity. *Journal of Fish Biology*, 37, 473-488.
- Paulay, G., and C. Meyer (2006). Dispersal and divergence across the greatest ocean region: do larvae matter? *Integr. Compar. Biol.*, 46, 269–281.
- Perron, F., and A. Kohn (1985). Larval dispersal and geographic distribution in coral reef gastropods of the genus *Conus*. *Proc. Fifth Internat. Coral Reef Congr.*, Tahiti, 4, 95–100.
- Pfenninger, M. (2004). Comparative analysis of range sizes in Helicidae s.l. (Pulmonata, Gastropoda). *Evol. Ecol. Res.*, 6, 359–376.
- Pielou, E. (1977). The latitudinal spans of seaweed species and their patterns of overlap. *Journal of Biogeography* 4, 299-311.
- Pielou, E. (1978). Latitudinal overlap of seaweed species: evidence for quasi-sympatric speciation. *Journal of Biogeography* 5, 227-238.
- Polechová J. and D. Storch (2008). Ecological niche. In: *Encyclopedia of Ecology*, Eds. Jorgensen and Fath. Elsevier, Oxford, 1088-1097.
- Pollock, K., Nichols, J., Brownie, C., and J. Hines (1990). Statistical inference for capture-recapture experiments. *Wildlife Monographs* 107, 1-98.

- Poulin, R., Luque, J., Guilhaumon, F., and D. Mouillot (2008) Species abundance distributions and numerical dominance in gastrointestinal helminth communities of fish hosts. *Journal of Helminthology* 82, 193–202.
- Predavec, M., Krebs, C., Danell, K., and R. Hyndman (2001). Cycles and synchrony in the Collared Lemming (*Dicrostonyx groenlandicus*) in Arctic North America. *Oecologia* 126:216–224
- Press, A. (1982). A comparison of the ecology of populations of *Rattus fuscipes* (Waterhouse) living in two distinct habitats. PhD Thesis, School of Biological Sciences, University of Sydney, NSW, Australia.
- Primack, R. (1980). Phenotypic variation of rare and widespread species of *Plantago*. Symposium on Rare and Endangered Species in New England, Cambridge, Mass. (USA).
- Pyke, G. (1982). Local geographic distributions of bumblebees near Crested Butte, Colorado: competition and community structure. *Ecology* 63:555–573.
- QLD Museum (1976). Fauna of eastern Australian rainforests I and II. Queensland Museum , Brisbane, QLD.
- Queensland. Nature Conservation Act 1992. Queensland Government printer, Brisbane, Queensland, Australia.
- Rabinowitz, D. (1981). Seven forms of rarity. In: H. Synge (ed). The biological aspects of rare plant conservation. Wiley, New York, 205-217.
- Rabinowitz, D., Cairns, S., and T. Dillon (1986). Seven forms of rarity and their frequency in the flora of the British Isles. In: (ed) M. Soule. Conservation Biology: the science of scarcity and diversity. Sinauer Associates, Sunderland, MA, 182-204.
- Raven, P., Johnson, G., Losos, J., Mason, K., and S. Singer (2008). Biology: 8th edition. McGraw Hill, New York, NY.
- Read, V. (1984). The survey of small mammal communities in forest environments of Australia. Masters Thesis, ANU, Canberra, ACT.

- Reaka, M. (1980). Geographic range, life history patterns, and body size in a guild of coral-dwelling mantis-shrimps. *Evolution*, 34, 1019-1030.
- Reed, J. (1992). A system for ranking conservation priorities for Neotropical migrant birds based on relative susceptibility to extinction. In: *Ecology and conservation of Neotropical migrant landbirds* (Eds) J. Hagan III and D. Johnston, Smithsonian Institute Press, Washington, 524-536.
- Resh, V., Bêche, L., and E. McElravy (2005). How common are rare taxa in long-term benthic macroinvertebrate surveys? *Journal of the North American Benthological Society*: 24, 976–989.
- Reveal, J. (1981). The concepts of rarity and population threats in plant communities, In: *Rare Plant Conservation*. (eds) Morse and Henefin. The New York Botanical Garden, Bronx, 41-46.
- Ritchie, M., and D. Tilman (1992). Interspecific competition among grasshoppers and their effect on plant abundance in experimental field environments. *Oecologia* 89: 524-532.
- Roberts, C. (2007). Scent marking. In Wolff, J., and Sherman (eds). *Rodent Societies: An ecological and Evolutionary perspective*. The University of Chicago Press, Chicago.
- Rogers, T. (1984). Rarity and chaos. *Mathematical Biosciences*, 72, 13-17.
- Rohde, K. (1992) Latitudinal gradients in species diversity: the search for the primary cause. *Oikos* 65: 514-527.
- Rohde, K. (1998) Latitudinal gradients in species diversity: area matters, but how much? *Oikos* 82: 184-190.
- Rohde, K. (2005). *Nonequilibrium Ecology*, Cambridge University Press, Cambridge.
- Rosenzweig, M. (1966). Community structure in sympatric Carnivora. *J. Mammal.* 47(4): 602–612.
- Room, P., Harley, K., Forno, I., and D. Sands (1981). Successful biological control of the floating weed salvinia. *Nature*, 294, 78-80.

- Rosenzweig, M. (1995). *Species Diversity in Space and Time*. Cambridge University Press, Cambridge.
- Rosenzweig, M., and J. Winakur (1969). Population ecology of desert rodent communities: habitat and environmental complexity. *Ecology*, 50, 558-572.
- Rosenzweig, M., and M. Lomolino (1997). Who gets the short bits of the broken stick? In: W. Kunin and K. Gaston (eds), *The Biology of rarity, Population and Community Biology Series*, Chapman and Hall, London, 63-86.
- Routledge, R. (1980). Bias in estimating the diversity of large, uncensused communities. *Ecology* 61:276–281.
- Roze, U. (1989). *North American Porcupine*. Smithsonian Institution Press, Wash. D.C.
- Roze, U. (2003). Porcupine *in* “Wild Mammals of North America: Biology, Management and Conservation” 2nd ed. Johns Hopkins Univ. Press, Baltimore, MD. 371-380.
- Rümmler, H. (1938). Neue Muriden aus Neuguinea. *Zeitschrift für Säugetierkunde* 10, 105-118.
- Savage, V., Gillooly J., Brown, J., West, G, and E. Charnov (2004). Effects of body size and temperature on population growth. *American Naturalist*, 163, 429-441
- Scheltema, R. (1989). Planktonic and non-planktonic development among prosobranch gastropods and its relationship to the geographic range of species. In: *Reproduction, Genetics, and Distributions of Marine Organisms* (eds Ryland, J.S. & Tyler, P.A.). Olsen & Olsen, Fredensborg, Denmark. 183–188.
- Schluter, D. (2001). *The ecology of adaptive radiation*. Oxford: Oxford University Press.
- Schneider K., Migge S., Norton, R., Scheu, S., Langel, R., Reineking, A., and M. Maraun (2004) Trophic niche differentiation in oribatid mites (Oribatida, Acari): evidence from stable isotope ratios. *Soil Biol. Biochem.* 36:1769–1774
- Schoener, T. (1987). The geographical distribution of rarity. *Oecologia*, 74, 161-173.

- Schwarz, C., and G. Seber (1999). Estimating animal abundance: a review III. *Statistical Science*, 14, 427-456.
- Schwarz, D., and S. Heckman (1980). The social role of scent marking in beaver (*Castor canadensis*). *Journal of Chemical Ecology*.
- Seaby, R., and Henderson, P. (2006). *Simply Tagging*. Pisces Conservation Ltd, IRC House, Lymington, England.
- Seaby, R., and P. Henderson (2006). *Species Diversity and Richness IV*. Pisces Conservation Ltd, IRC House, Lymington, England.
- Seber, G. (1982). *The estimation of animal abundance and related parameters*. Griffin, London.
- Seber, G. A. F. (1986). A review of estimating animal abundance. *Biometrics*, 42, 267-292.
- Shkedy, Y., and U. Safriel (1992). Niche breadth of two lark species in the desert and the size of their geographical ranges. *Ornis Scandinavica* 23, 89-95.
- Sih, A. (1982). Foraging Strategies and the Avoidance of Predation by an Aquatic Insect, *Notonecta Hoffmanni*. *Ecology*, Vol. 63, 786-796.
- Singleton, G. (2008). House Mouse *Mus musculus*. In: S. Van Dyck and R. Strahan (eds), *The Mammals of Australia*, Reed New Holland, Australasia.
- Smith, G. (1985). Population and Habitat Selection of the Noisy Scrub-bird, *Atrichornis clamosus*, 1962-83. *Australian Wildlife Research* 12, 479-485.
- Söderström, O. (1989) Effects of temperature and food quality on life-history parameters in *Parameletus chelifera* and *P. minor* (Ephemeroptera): a laboratory study. *Freshwater Biology* 20:3, 295-303.
- Solow, A. (1993). A simple test for change in community structure. *J. Anim. Ecol.*, 62(1), 191-193.
- Spitzer, K., and J. Lepš (1988). Determinants of temporal variation in moth abundance. *Oikos*, 53, 31-36.



- Stanton, P., and D. Stanton (2006). Vegetation mapping of the Wet Tropics Heritage Area. Wet Tropics Management Authority, Cairns, Australia.
- Stevens, G. (1989). The latitudinal gradient in geographic range: how so many species coexist in the tropics. *American Naturalist*, 133, 240-256.
- Stirling, G., and Wilsey, B. (2001). Empirical Relationships between Species Richness, Evenness, and Proportional Diversity. *Am. Nat.* 2001. Vol. 158, 286–299.
- Sreatfeild, C. (2009). The effects of habitat fragmentation on the demography and population genetic structure of *Uromys caudimaculatus*. PhD Thesis, Queensland University of Technology, Brisbane, Australia.
- Strong, D. (1992). Are trophic cascades all wet? Differentiation and donor-control in speciose ecosystems. *Ecology* 73: 747-754.
- Sutherland, W., and S. Baillie (1993). Patterns in the distribution, abundance, and variation of bird populations. *Ibis*, 135, 209-210.
- Taber, A., Novaro, A., Neris, N., and F. Colman (1997). The food habits of sympatric jaguar and puma in the Paraguayan Chaco. *Biotropica*, 29, 204–213.
- Tate, G. (1951). Results of the Archbold Expedition. No. 65. The rodents of Australia and New Guinea. *Bulletin of the American Museum of Natural History*, Vol. 97, 183-430.
- Taylor, J. (1961). Reproductive biology of the Australian Bush Rat (*Rattus assimilis*). University of California Publication, 60.
- Taylor, J., and B. Horner (1970). Reproduction in the mosaic-tailed rat *Melomys cervinipes* (Rodentia: Muridae), *Australian Journal of Zoology*, Vol. 18, 171-184.
- Theimer, T. (2001) Seed scatterhoarding by white-tailed rats: consequences for seedling recruitment by an Australian rain forest tree. *Journal of Tropical Ecology*, 17, 177-189.
- Theimer, T. (2003). Intraspecific variation in seed size affects scatterhoarding behaviour of an Australian tropical rain-forest rodent *Journal of Tropical Ecology* 19:95–98.

- Thompson, K., Hodgson, J., and K.Gaston (1998). Abundance-Range Size Relationships in the Herbaceous Flora of Central England. *J. Ecol.*, 86, 439-448.
- Thompson, K., Gaston, K., and S. Band. (1999). Range size, dispersal and niche breadth in the herbaceous flora of central England. *J. Ecol.*, 87, 150–155.
- Thornton, D., Sunkist, M. and B. Martin (2004). Ecological separation within newly sympatric populations of coyotes and bobcats in south-central Florida Source. *Journal of Mammalogy*. 85(5), 973-982.
- Tilman, D. (1982). Resource competition and community structure. Princeton University Press.
- Tilman, D. (1988). Plant strategies and the dynamics and structure of plant communities. Princeton University Press.
- Tilman, D. (1990). Constraints and tradeoffs: toward a predictive theory of competition and succession. *Oikos* 58: 3-15.
- Tilman, D. (1996). Biodiversity: Population Versus Ecosystem Stability. *Ecology*, Vol. 77, No. 2, 350-363.
- Torgersen, T., Luly, J., DeDeckker, P., Jomes, M., Searle, D., Chivas, A., and W. Ullman(1988). Late Quaternary environments of the Carpentaria Basin, Australia. *Paleogeography, Paleoclimatology, Paleoecology*, Vol. 67, 445-461.
- Tracey, J. (1982). The vegetation of the humid tropical region of north Queensland. CSIRO, Melbourne.
- Turin, H., and P. den Boer (1988). Changes in the distribution of carabid beetles in The Netherlands since 1880. II. Isolation of habitats and long-term time trends in the occurrence of carabid species with different powers of dispersal (Coleoptera, Carabidae). *Biological Conservation* 44, 179-200.
- Udvardy, M. (1959). Notes on the ecological concepts of habitat, biotope, and niche. *Ecology* 40, 725-728.

- Unwin, G., Applegate, G., Stocker, G., and D. Nicholson (1988) Initial effects of tropical cyclone 'Winifred' on forests in north Queensland. *Proc. Ecol. Soc. Aust.* 15, 283–96.
- Urban, M. (2004) Disturbance heterogeneity determines freshwater metacommunity structure. *Ecology* 85:11, 2971-2978.
- Vallade, M. and B. Houchmandzadeh (2003). Analytical solution of a neutral model of biodiversity. *Phys. Rev. E*, 68, 061902.
- van Oosterzee, P. (1997). *Where worlds collide: The Wallace Line*. Reed Books, Sydney.
- van Oosterzee, P. (2006). Drawing the Wallace Line. In: (Eds) Merrick, Archer, Hickey, Lees, *Evolution and biogeography of Australian Vertebrates*, Auscipub Pty Ltd, Australia.
- Vandermeer, J. (1982). To be rare is to be chaotic. *Ecology*, 63, 1167-1168.
- Vanschoenwinkel B., De Vries, C., Seaman, M., L. Brendonck (2007). The role of metacommunity processes in shaping invertebrate rock pool communities along a dispersal gradient. *Oikos* 116:1255–1266.
- Vartanyan, S., Garutt, V., and A. Sher. (1993). Holocene dwarf mammoths from Wrangel Island in the Siberian Arctic. *Nature*, 362, 337-340.
- Volkov, I., Banavar, J., Hubbell, S., and A. Maritan (2003). Neutral theory and relative species abundance in ecology. *Nature*, 424, 1035–1037.
- Volterra, V.(1926). Fluctuations in the abundance of a species considered mathematically. *Nature* 118: 558-60.
- Volterra, V. (1931). Variations and fluctuations of the number of individuals in animal species living together. In: *Animal Ecology*. McGraw–Hill, 1931. Translated from 1928 edition by R. N. Chapman.

- Voznessenskaya, V., Parfyonova, V., and E. Zinkevich (1992). Individual odour-types. In: R. Doty and D. Muller-Scha Marmotinirze (eds), Chemical signals in vertebrates VI, 503-508. New York: Plenum.
- Walker, J. (2006). Resource use and rarity among frugivorous birds in a tropical rain forest on Sulawesi. *Biological Conservation*, 130, 60–69
- Warneke, R. (1971). Field study of the bush rat (*Rattus fuscipes*). *Wildlife Control Victoria* No.14.
- Watts, A. (1997). Population Ecology and Reproductive Seasonality in Three Species of *Antechinus* (Marsupialia:Dasyuridae) in the Wet Tropics of Queensland. *Wildlife Research*, 1997.
- Watts, C. (1977). The foods eaten by some Australian rodents (Muridae). *Aust. Wildl. Res.* 4, 151-158.
- Watts, C., and H. Aslin (1981). *Rodents of Australia*. Angus and Robertson, Australia.
- Watts, C., Baverstock, P., Birrell, J. and Kreig, M. (1992). Phylogeny of the Australian rodents (Muridae): a molecular approach using microcomplement fixation of albumin. *Aust. J. Zool.* 40, 81-90.
- Watts, C. and Baverstock, P. (1994a). Evolution in New Guinea Muridae (Rodentia) assessed by microcomplement fixation of albumin. *Aust. J. Zool.* 42, 295-306.
- Watts, C. and Baverstock, P. (1994b). Evolution in some South-east Asian Murinae (Rodentia), as assessed by microcomplement fixation of albumin, and their relationship to Australian murines. *Aust. J. Zool.*, 42, 711-722.
- Watts, C. and Baverstock, P. (1995a). Evolution in the Murinae (Rodentia) assessed by microcomplement fixation of albumin. *Aust. J. Zool.*, 43, 105-118.
- Watts, C. and Baverstock, P. (1995b). Evolution in some African Murinae (Rodentia) assessed by microcomplement fixation of albumin. *Journal of African Zoology.*, 109, 423-433.


- Webb, S. (1984). On two kinds of faunal turnover. In: W. Berggren and J. Van Couvering (eds). *Catastrophes and earth history: the new uniformitarianism*. Princeton Press, Princeton, NJ, 417-436.
- Watts, C., and P. Baverstock (1994). Evolution in New Guinea Muridae (Rodentia) assessed by microcomplement fixation of albumin. *Australian Journal of Zoology*, Vol. 42, 295-306.
- Watts, C., and Baverstock, P. (1996). Phylogeny and biogeography of some Indo-Australian murid rodents. 47-50. In Kitchener, D., and Suyanta, A. (eds). *Proceedings of the First International Conference on Eastern Indonesian-Australian Vertebrate Fauna*, Manado, Indonesia, November 22-26, Western Australian Museum, Perth.
- Wellesley-Whitehouse, H. (1981). A study on the comparative ecologies of two rainforest rodents, *Uromys caudimaculatus* and *Rattus fuscipes coracius* in north Queensland. Honours Thesis, Department of Zoology, James Cook University, Townsville, Queensland, Australia.
- Wellington, G., and B. Victor (1989). Planktonic larval duration of one hundred species of Pacific and Atlantic damselfishes (Pomacentridae). *Mar. Biol.*, 101, 557–568.
- Werner, E., Gilliam, J., Hall, D., and G. Mittelbach (1983). An experimental test of the effects of predation risk on habitat use in fish. *Ecology* 64:1540–1548.
- Werner, E., and B. Anholt (1993). Ecological consequences of the trade-off between growth and mortality rates mediated by foraging activity. *Am. Nat.* 142, 242.
- Werth, S., Wagner, H., Gugerli, F., Holderegger, R., Csencsics, Kalwij, D., and M. Jesse M. (2006). Quantifying dispersal and establishment limitation in a population of an epiphytic lichen. *Ecology*, 87(8), 2006, 2037–2046.
- White, G. (1983). Numerical estimation of survival rates from band recovery and biotelemetry data. *J. Wildl. Manage.* 47, 716-728.

- White, G., Anderson, D., Burnham, K., and D. Otis (1982). Capture-recapture and removal methods for sampling closed populations. Los Alamos National Laboratory, Los Alamos, New Mexico.
- White, M. (2006). Environments of the geological past. In: Merrick, J., Archer, M., Hickey, G., and Lee, M. (eds), Evolution and Biogeography of Australasian Vertebrates. Auscipub Pty Ltd. NSW.
- Whitfield, J. (2002). Neutrality versus the niche. *Nature*. 417: 480-481.
- Whittaker, R. (1972). Evolution and measurement of species diversity. *Taxon*, 21, 213-251.
- Whittaker, R., Levin, S., and Root, R. (1973). Niche, habitat, and ecotype. *Am. Nat.* 107: 3212-3338.
- Whittaker, R., and Levins, S. (1975). Niche: theory and application. Stroudsburg, Wiley.
- Williams, B., Nichols, J., and M. Conroy (2002). Analysis and management of animal populations. Academic Press, London.
- Williams, S. (2006). Vertebrates of the Wet Tropics rainforests of Australia: Distribution and Biodiversity. Cooperative Research Centre for Tropical Rainforest Ecology and Management, Cairns, Queensland, Australia.
- Williams, S., and H.Marsh (1998) Changes in small mammal assemblage structure across a rain forest/open forest ecotone. *Journal of Tropical Ecology* 14:187–198.
- Wilson, K., and D. Anderson (1985). Evaluation of a nested grid approach for estimating density. *J. Wildl. Manage.* 49, 675-678.
- Winter, J. (1984). The Thornton Peak Melomys, *Melomys hadrourus* (Rodentia: Muridae): a new rainforest species from northeastern Queensland, Australia. *Memoirs of the Queensland Museum* 21, 519-539.

- Winter, J., Whitford, D., and R. Rader (2008) Tree Mouse *Pogonomys* sp. In: (eds) Van Dyck and Strahan, The mammals of Australia. Reed New Holland, Sydney, Australia,
- Wood, D. (1971). The ecology of *Rattus fuscipes* and *Melomys cervinipes* in a south-east Queensland rainforest. *Aust. J. Zool.* 19: 371-392.
- Woolnough, A., Johnson, C., and A. Horsup (1998). The short-term effect of radio-packages on a free-ranging large herbivore, the northern hairy-nosed wombat. *Wildlife Research* 25 (5): 561-565.
- Zar, J.H. 1999. *Biostatistical analysis*. 4th edition. Prentice Hall, Upper Saddle River, NJ.


APPENDIX 1

Pes morphology of north Queensland rodents (\* IDP = Interdigital pads)






*Melomys cervinipes* - Adult female




Pes: 27.0 mm (without claws) ; 28.0 mm (with claws)

*Uromys hadrourus* - Adult female




Pes: 39.5.0 mm (without claws) ; 41.50 mm (with claws)

*Melomys burtoni* - Adult male



Pes: 26.2 mm (without claws) ; 27.2 mm (with claws)

*Rattus fuscipes* - Adult female



Pes: 34.0 mm (without claws) ; 35.0 mm (with claws)